

TABLE A. INDIVIDUAL LOCATION DATA AT NORTH CAROLINA FOR GROUP IV CONVENTIONAL AND LL SOYBEAN - 2013.

BRAND VARIETY OR VARIETY	EP WASHINGTON BU/A	EP LENOIR BU/A	LP LENOIR [¶] BU/A	AVERAGE BU/A
Stine 49LD02	75.8**	70.2**	59.4*	68.5**
Progeny P4930LL	68.2	69.6*	59.6**	65.8*
SH 4913LL	69.3	67.4*	53.8*	63.5*
HBK LL4953	72.3*	67.2*	47.6	62.4
HALO 5:01LL	72.0*	64.0	48.7	61.6
AG 4730	65.0	67.4*	50.0	60.8
+HALO X496LL	70.4*	65.7*	45.1	60.4
+S09-9943	65.5	55.0	56.3*	58.9
+S09-10871	71.4*	54.4	50.3	58.7
HBK LL4950	62.5	66.1*	46.5	58.4
Progeny 4928LL	66.1	56.9	50.3	57.8
USG 74G82L	62.8	64.3	46.1	57.7
+S09-13608	68.3	56.4	43.1	56.0
HALO 4:94LL	63.3	59.0	45.2	55.8
Progeny 4819LL	62.6	60.0	44.0	55.5
SH 4714LL/STS	62.4	60.3	40.7	54.5
HALO 4:97LL/STS	58.8	60.5	40.4	53.2
HALO 4:95LL	61.9	48.8	42.1	50.9
Progeny P4560LL	62.1	48.5	40.2	50.3
HBK LL4850	63.7	55.5	30.1	49.8
HBK LL4650	50.7	47.5	35.4	44.5
+HALO X466LL	52.9	38.5	38.0	43.1
HBK LL4653	55.2	42.0	25.8	41.0
MEAN	64.5	58.5	45.2	56.0
ADJ R2	62.8	78.8	80.6	89.9
BLSD (K-50)	6.2	5.3	7.4	6.0
C.V.	9.5	9.6	14.5	7.9
S.E.	2.7	2.5	3.3	1.2
ERROR D.F.	89	84	61	44

+EXPERIMENTAL

**HIGHEST YIELDER. *NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELDER.

[¶] Refer to SEASONAL CONDITIONS section

EP = Early Planted

LP = Late Planted