

TABLE B. INDIVIDUAL LOCATION DATA AT NORTH CAROLINA FOR GROUP V EARLY (5.0 - 5.5)
CONVENTIONAL AND LL SOYBEAN - 2013.

BRAND VARIETY OR VARIETY	EP WASHINGTON BU/A	EP LENOIR BU/A	LP LENOIR [†] BU/A	LP ROWAN BU/A	AVERAGE BU/A
+N02-7002	73.7*	66.1*	56.3*	42.4	59.6**
+JTN-5303	68.8*	65.6*	61.9**	40.9	59.3*
JTN-5110	66.9*	66.3*	59.3*	44.6*	59.3*
SH 5512LL	73.5*	63.9*	53.7*	43.4	58.6*
Stine 54LE23	72.0*	66.8*	47.3	47.3*	58.3*
GoSoy 5010LL	67.4*	64.5*	52.6	48.6**	58.3*
JTN-5503	64.5	69.9*	61.1*	36.7	58.1*
Osage	75.7**	64.6*	45.9	45.7*	58.0*
HALO 5:45LL	65.8*	61.2	59.4*	45.1*	57.9*
HALO 5:01-5LL	63.0	69.2*	53.1*	43.6	57.2*
Jake	71.2*	63.4*	52.3	41.8	57.2*
Ozark	73.4*	63.3	45.4	45.9*	57.0*
UA 5213C	70.5*	64.3*	45.4	46.5*	56.7*
SS LL513N	70.2*	61.6	48.5	45.3*	56.4*
SSLL 511N	68.7*	62.6	44.4	46.5*	55.6*
GoSoy 5312LL	61.5	67.0*	46.3	47.0*	55.5*
HALO 5:26LL	70.1*	63.6*	41.4	46.6*	55.4*
+JTN-4307	67.4*	65.6*	41.3	43.9	54.5*
+NCC07-7506	69.6*	61.9	43.8	42.9	54.5*
Fowler	64.2	70.7**	40.1	40.6	53.9*
Progeny 5160LL	68.1*	62.7	39.9	43.4	53.5*
Glenn	60.4	57.2	46.6	47.2*	52.8*
+S09-13635	57.0	65.2*	44.1	41.8	52.0*
AGS 553LL	60.8	59.8	44.4	41.9	51.7
+HALO X530LL/STS	67.7*	54.5	43.7	40.6	51.6
JTN-5203	61.1	47.4	60.5*	37.1	51.5
Stine 54LD00	53.7	67.4*	40.2	39.9	50.3
Progeny 5460LL	60.7	50.6	43.8	45.8*	50.2
SH 5212LL	61.7	60.0	38.1	40.8	50.1
USG 7553nRS	63.0	59.0	40.5	37.6	50.0
GoSoy 5410LL	58.0	56.8	40.0	45.0*	49.9
+N08-93	61.0	58.9	35.9	36.8	48.2
Hutcheson	62.4	63.0	32.9	32.8	47.8
HBK LL5350	64.6	53.1	33.5	39.1	47.6
MEAN	65.8	62.3	46.6	42.8	54.4
ADJ R2	55.2	53.9	67.6	71.1	85.3
BLSD (K-50)	10.8	7.4	9.1	4.4	7.7
C.V.	12.4	9.4	16.3	9.7	9.4
S.E.	3.6	2.9	3.8	1.8	1.2
ERROR D.F.	128	97	94	130	99

+EXPERIMENTAL

**HIGHEST YIELDER. *NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELDER.

[†] Refer to SEASONAL CONDITIONS section

EP = Early Planted; LP = Late Planted