

TABLE C. INDIVIDUAL LOCATION DATA AT NORTH CAROLINA FOR GROUP V LATE (5.6 - 5.9)
CONVENTIONAL AND LL SOYBEAN - 2013.

| BRAND VARIETY OR VARIETY | EP WASHINGTON BU/A | EP LENOIR BU/A | LP LENOIR [¶] BU/A | LP ROWAN BU/A | AVERAGE BU/A |
|-----------------------------|--------------------------|----------------------|-----------------------------------|---------------------|-----------------|
| UA 5612 | 75.9* | 65.7** | 39.1 | 44.0** | 56.2** |
| AGS 5911LL | 67.4 | 62.8* | 51.6* | 40.4 | 55.6* |
| Fowler | 80.3* | 61.5* | 46.3 | 34.1 | 55.5* |
| Jake | 67.9 | 63.9* | 50.9* | 38.2 | 55.3* |
| Osage | 81.3** | 54.6 | 47.5* | 34.2 | 54.4* |
| SS LL563NSTS | 73.7* | 55.5 | 45.9 | 38.2 | 53.3* |
| SH 5614LL/STS | 69.9 | 60.3* | 46.2 | 36.5 | 53.2* |
| +N08-174 | 75.4* | 60.6* | 39.1 | 37.5 | 53.1* |
| NC Miller | 73.3* | 62.8* | 40.4 | 35.8 | 53.1* |
| USG 75J90R | 75.7* | 55.7 | 41.3 | 39.5 | 53.1* |
| Progeny 5960LL | 65.8 | 58.3 | 54.3** | 33.6 | 53.0* |
| SH 5912LL | 64.3 | 61.3* | 50.3* | 33.4 | 52.4* |
| SS LL595N | 62.8 | 62.2* | 46.8* | 33.8 | 51.4* |
| Stine 58LC23 | 70.3* | 60.3* | 38.5 | 35.6 | 51.2* |
| Hutcheson | 74.9* | 55.7 | 34.8 | 35.6 | 50.2* |
| MEAN | 71.9 | 60.1 | 44.9 | 36.7 | 53.4 |
| ADJ R2 | 51.7 | 58.3 | 73.3 | 81.0 | 91.7 |
| BLSD (K-50) | 11.2 | 6.4 | 7.9 | 3.4 | 11.7 |
| C.V. | 11.7 | 8.0 | 13.8 | 8.6 | 9.2 |
| S.E. | 3.8 | 2.2 | 3.1 | 1.4 | 1.1 |
| ERROR D.F. | 52 | 52 | 41 | 52 | 42 |

+EXPERIMENTAL

**HIGHEST YIELDER. *NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELDER.

[¶] Refer to SEASONAL CONDITIONS section

EP = Early Planted

LP = Late Planted