

TABLE D. INDIVIDUAL LOCATION DATA AT NORTH CAROLINA FOR GROUP VI CONVENTIONAL AND LL SOYBEAN - 2013.

BRAND VARIETY OR VARIETY	EP WASHINGTON BU/A	EP LENOIR BU/A	LP LENOIR [¶] BU/A	LP ROWAN BU/A	AVERAGE BU/A
+N08-145	80.9**	63.1**	37.5*	45.2**	56.7**
+NCC07-8138	78.1*	60.5*	35.0*	43.4*	54.2*
+N07-187	77.4*	61.4*	32.5	42.2*	53.4*
NC Roy	77.5*	60.1*	32.6	42.1	53.1*
USG 620nRR	72.3	62.4*	38.5**	38.2	52.9*
+N05-7375	66.4	56.5	38.0*	37.5	49.6*
+N06-7023	73.6	55.2	27.3	41.5	49.4*
USG 76G10L	70.7	46.8	36.8*	41.6	48.9*
+N05-7353	74.4	48.8	28.3	36.7	47.1
+NCC06-1090	78.7*	26.6	10.9	27.0	35.8
+NCC07-7961	56.7	10.1	5.0	21.8	23.4
MEAN	73.3	50.1	29.3	37.9	47.7
ADJ R2	72.3	93.2	92.2	90.0	92.1
BLSD (K-50)	5.6	5.2	5.1	3.1	7.9
C.V.	7.5	10.9	15.5	8.5	14.1
S.E.	2.5	2.4	2.3	1.4	1.5
ERROR D.F.	42	38	27	36	30

+EXPERIMENTAL

**HIGHEST YIELDER. *NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELDER.

[¶] Refer to SEASONAL CONDITIONS section

EP = Early Planted

LP = Late Planted