

TABLE E. INDIVIDUAL LOCATION DATA AT NORTH CAROLINA FOR GROUP VII-VII CONVENTIONAL SOYBEAN - 2013.

BRAND VARIETY OR VARIETY	EP	EP	LP	LP	AVERAGE BU/A
	WASHINGTON BU/A	LENOIR BU/A	LENOIR [†] BU/A	ROWAN BU/A	
N 7003CN	64.7*	71.4**	48.3**	40.7	56.3**
AGS Woodruff	67.4*	66.2	35.3	46.0**	53.7*
+N08-391	66.6*	64.9	34.6	45.8*	53.0*
+N05-7462	70.0**	59.5	34.2	43.5*	51.8*
+N06-7564	66.5*	56.9	36.8	40.2	50.1
USG 7732nRR	60.0	58.9	31.6	41.4	48.0
+N08-447	57.9	61.4	30.2	42.1	47.9
+N05-7396	61.3	56.7	30.9	41.3	47.5
+N08-374	68.0*	43.6	35.8	42.4	47.4
N8001	62.5*	52.4	32.8	38.1	46.4
+N05-7432	55.4	53.4	36.9	37.4	45.8
NC Raleigh	55.3	60.8	30.0	36.1	45.6
N7002	54.4	55.8	34.2	36.4	45.2
+N05-7452	57.7	51.7	27.6	37.9	43.7
+NCC06-899	56.7	48.5	17.6	37.3	40.0
MEAN	61.6	57.5	33.1	40.4	48.2
ADJ R2	59.4	83.6	63.3	74.4	91.8
BLSD (K-50)	8.3	4.8	8.5	3.6	5.7
C.V.	11.2	8.2	22.5	8.3	9.1
S.E.	3.1	2.1	3.3	1.5	1.0
ERROR D.F.	52	52	55	56	42

+EXPERIMENTAL

**HIGHEST YIELDER. *NOT SIGNIFICANTLY DIFFERENT FROM HIGHEST YIELDER.

[†] Refer to SEASONAL CONDITIONS section

EP = Early Planted

LP = Late Planted